

## Asymptotic Representation of Solutions of Equation

$$\dot{y}(t) = \beta(t)[y(t) - y(t - \tau(t))]$$

Josef Diblík\*

*Department of Mathematics. Faculty of Electrical Engineering and Computer Science.*

metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

Received September 11, 1996

In this paper the equation  $\dot{y}(t) = \beta(t)[y(t) - y(t - \tau(t))]$  is considered where  $\tau \in C(I_{-1}, \mathbb{R}^+)$ ,  $I_{-1} = [t_{-1}, \infty)$ ,  $\mathbb{R}^+ = (0, \infty)$ ,  $t - \tau(t)$  is an increasing function on  $I_{-1}$ ,  $\tau(t) \leq \tilde{\tau}$ ,  $t \in I_{-1}$ ,  $0 < \tilde{\tau} = \text{const}$  and  $\beta \in C(I_{-1}, \mathbb{R}^+)$ . Results about asymptotic structure and behavior of the solutions of this equation are proved. A connection with the equation  $\dot{x}(t) = -c(t)x(t - \tau(t))$  and the asymptotic representation of its solutions is obtained. © 1998 Academic Press

### 1. INTRODUCTION

In this paper, we consider the equation

$$\dot{y}(t) = \beta(t)[y(t) - y(t - \tau(t))], \quad (1)$$

where  $\tau \in C(I_{-1}, \mathbb{R}^+)$ ,  $I_{-1} = [t_{-1}, \infty)$ ,  $t_{-1} \in \mathbb{R}$ ,  $\mathbb{R}^+ = (0, \infty)$ ,  $t - \tau(t)$  is an increasing function on  $I_{-1}$ ,  $\tau(t) \leq \tilde{\tau}$ ,  $t \in I_{-1}$ ,  $0 < \tilde{\tau} = \text{const}$ , and  $\beta \in C(I_{-1}, \mathbb{R}^+)$ , although in many affirmations,  $\tau \in C(I, \mathbb{R}^+)$ ,  $\beta \in C(I, \mathbb{R}^+)$ ,  $I = [t_0, \infty)$ ,  $t_{-1} = t_0 - \tau(t_0)$  is sufficient.

We prove results about asymptotic structure and behavior of the solutions of (1). A connection with the equation

$$\dot{x}(t) = -c(t)x(t - \tau(t)) \quad (2)$$

is shown and, as application of the obtained results, asymptotic representation of its solution is obtained.

\* E-mail: <diblik@dmf.fee.vutbr.cz>.

Equation (1) was investigated, e.g., in papers by F. V. Atkinson and J. R. Haddock [2] and S. N. Zhang [18]. Some questions related to the preceding problems are discussed, e.g., in the works by O. Arino, I. Györi, and M. Pituk, [1], R. Bellman and K. L. Cooke [3], K. Gopalsamy [9], I. Györi and M. Pituk [7, 8], E. Kozakiewicz [11] and T. Krisztin [12]. Therefore some comparisons to known results are made in the text of the paper.

## 2. PRELIMINARY

Consider the system of functional differential equations of retarded type,

$$\dot{y}(t) = f(t, y_t), \quad (3)$$

where  $y \in \mathbb{R}^n$ ,  $y_t$  is an element from the space of continuous functions  $C = C([-\tau, 0], \mathbb{R}^n)$ ,  $y_t(\theta) = y(t + \theta)$  where  $\theta \in [-\tau, 0]$ ,  $f: \Omega \mapsto \mathbb{R}^n$ ,  $\Omega$  is an open subset of  $\mathbb{R} \times C$  and  $f$  is a continuous mapping such that the element  $(\delta, \pi) \in \Omega$  determines a unique solution  $y(\delta, \pi)$  on its maximal existence interval  $D_{\delta, \pi} = [\delta, a)$ ,  $\delta < a \leq \infty$  (see, e.g., [10]). In the case of the linear equation (1) these conditions are satisfied and  $a = \infty$ .

In the paper, we use the topological principle of T. Ważewski [17] in the form of K. P. Rybakowski [16] for retarded functional differential equations. This modification applies the ideas of B. S. Razumikhin (see, e.g., [15]). Now we give a concept of this principle. (However, as usual, if  $\omega \subset \mathbb{R} \times \mathbb{R}^n$ , then  $\text{int } \omega$  and  $\partial\omega$  denote the interior and the boundary of  $\omega$ , respectively.)

**DEFINITION 1** [16]. Let the continuously differentiable functions  $l_i(t, y)$ ,  $i = 1, 2, \dots, p$  and  $m_j(t, y)$ ,  $j = 1, 2, \dots, q$ ,  $p^2 + q^2 > 0$  be defined on some open set  $\omega_0 \subset \mathbb{R} \times \mathbb{R}^n$ . The set,

$$\omega = \{(t, y) \in \omega_0 : l_i(t, y) < 0, m_j(t, y) < 0, \\ i = 1, 2, \dots, p, j = 1, 2, \dots, q\}$$

is called a *regular polyfacial set* with respect to the system (3) if  $(\alpha)$  to  $(\gamma)$  in the following text hold:

$(\alpha)$  For  $(t, \pi) \in \mathbb{R} \times C$  such that  $(t + \theta, \pi(\theta)) \in \omega$  for  $\theta \in [-\tau, 0]$ , we have  $(t, \pi) \in \Omega$ .

$(\beta)$  For all  $i = 1, 2, \dots, p$ , all  $(t, y) \in \partial\omega$  for which  $l_i(t, y) = 0$ , and all  $\pi \in C$  for which  $\pi(0) = y$  and  $(t + \theta, \pi(\theta)) \in \omega$ ,  $\theta \in [-\tau, 0]$  it fol-

lows that  $Dl_i > 0$  where,

$$Dl_i \equiv \frac{\partial l_i(t, y)}{\partial t} + \sum_{r=1}^n \frac{\partial l_i(t, y)}{\partial y_r} f_r(t, \pi).$$

( $\gamma$ ) For all  $j = 1, 2, \dots, q$ , all  $(t, y) \in \partial\omega$  for which  $m_j(t, y) = 0$ , and all  $\pi \in C$  for which  $\pi(0) = y$  and  $(t + \theta, \pi(\theta)) \in \omega$ ,  $\theta \in [-\tau, 0)$  it follows that  $Dm_j < 0$ .

The elements  $(t, \pi) \in \mathbb{R} \times C$  are in the sequel assumed to be such that  $(t, \pi) \in \Omega$ .

**DEFINITION 2.** A system of initial functions  $p_{A, \omega}$  with respect to the nonempty sets  $A$  and  $\omega$  where  $A \subset \bar{\omega} \subset \mathbb{R} \times \mathbb{R}^n$  is defined as a continuous mapping  $p: A \rightarrow C$  such that ( $\alpha$ ) and ( $\beta$ ) in the following text hold:

( $\alpha$ ) If  $z = (t, y) \in A \cap \text{int } \omega$  then  $(t + \theta, p(z)(\theta)) \in \omega$  for  $\theta \in [-\tau, 0]$ .

( $\beta$ ) If  $z = (t, y) \in A \cap \partial\omega$  then  $(t + \theta, p(z)(\theta)) \in \omega$  for  $\theta \in [-\tau, 0)$  and  $(t, p(z)(0)) = z$ .

The following lemma describes the main result of the paper by K. P. Rybakowski [16].

**LEMMA 1.** Let  $\omega \subset \omega_0$  be a regular polyfacial set with respect to the system (3) and let  $W$  be defined as follows,

$$W = \{(t, y) \in \partial\omega: m_j(t, y) < 0, j = 1, 2, \dots, q\}.$$

Let  $Z \subset W \cup \omega$  be a given set such that  $Z \cap W$  is a retract of  $W$  but not a retract of  $Z$ . Then for each fixed system of initial functions  $p_{Z, \omega}$  there is a point  $z_0 = (\sigma_0, y_0) \in Z \cap \omega$  such that for the corresponding solution  $y(\sigma_0, p(z_0))(t)$  of (3) we have  $(t, y(\sigma_0, p(z_0))(t)) \in \omega$  for each  $t \in D_{\sigma_0, p(z_0)}$ .

### 3. SOME PROPERTIES OF SOLUTIONS

The solution  $y = y(t)$  of (1) on interval  $I_{-1}$  is defined in a usual way as a function  $y(t) \in C(I_{-1}, \mathbb{R}) \cap C^1(I, \mathbb{R})$  which satisfies (1) on  $I$ . Denote, as  $y^\varphi(t)$ , a solution of (1) to be generated by an initial function  $\varphi(t)$ , which is defined and continuous on an initial interval  $[t_{-1}, t_0]$ .

We start from following several lemmas. Obviously in view of the form of (1) the following lemma holds:

**LEMMA 2.** If  $y(t) = y^\varphi(t)$  is a solution of (1) on  $I_{-1}$  generated by an initial function  $\varphi(t)$ , then the function  $Ky^\varphi(t) + L$  is a solution of (1) on  $I_{-1}$ , generated by initial function  $K\varphi(t) + L$  for every constant  $K, L \in \mathbb{R}$ .

LEMMA 3. Let the initial function  $\varphi(t)$  be defined and continuous on  $[t_{-1}, t_0]$  and

$$\varphi(t) < \varphi(t_0), \quad (4)$$

or

$$\varphi(t) > \varphi(t_0), \quad (5)$$

where  $t \in [t_{-1}, t_0)$ . Then the corresponding solution  $y^\varphi(t)$  is increasing in the case of inequality (4) or decreasing in the case of inequality (5) on  $I$ .

*Proof.* Because it follows from (1)  $\text{sign } \dot{y}^\varphi(t_0) = +1$  in the case (4) and  $\text{sign } \dot{y}^\varphi(t_0) = -1$  in the case (5). The case  $\dot{y}^\varphi(t^*) = 0$  for a  $t^* \in (t_0, \infty)$  and  $\text{sign } \dot{y}^\varphi(t) \neq 0$  where  $t \in (t_0, t^*)$  is impossible because, as follows from (1),  $y(t^*) \neq y(t^* - \tau(t^*))$ . The lemma is proved.

LEMMA 4. Let the initial function  $\varphi(t)$  be continuous increasing positive (or decreasing negative) on  $[t_{-1}, t_0]$ . Then

$$0 < |y^\varphi(t)| \leq |\varphi(t_0)| e^{\int_{t_0}^t \beta(s) ds}. \quad (6)$$

*Proof.* Let, e.g.,  $\varphi(t)$  be positive. By Lemma 3  $y^\varphi(t)$  increases on  $I$ . From representation,

$$y^\varphi(t) = e^{\int_{t_0}^t \beta(s) ds} \left[ \varphi(t_0) - \int_{t_0}^t \beta(s) y^\varphi(s - \tau(s)) e^{-\int_{t_0}^s \beta(q) dq} ds \right], \quad t \in I,$$

the inequality (6) follows. If  $\varphi(t)$  is negative, we proceed by analogy. The lemma is proved.

#### 4. REPRESENTATION OF SOLUTIONS

Consider the differential inequality,

$$\dot{\omega}(t) \leq \beta(t) [\omega(t) - \omega(t - \tau(t))], \quad (7)$$

where  $t \in I$ . The solution  $\omega = \omega(t)$  of inequality (7) on interval  $I_{-1} = [t_{-1}, \infty)$  is a function  $\omega(t) \in C(I_{-1}, \mathbb{R}) \cap C^1(I, \mathbb{R})$  which satisfies the inequality (7) on  $I$ .

THEOREM 1. If  $y(t)$  is a solution of (1) on  $I_{-1}$ , then there is a solution  $\omega(t)$  of inequality (7) on  $I_{-1}$  such that

$$y(t) > \omega(t), \quad t \in I_{-1}. \quad (8)$$

Conversely, if  $\omega(t)$  is a solution of inequality (7) on  $I_{-1}$ , then there is a solution  $y(t)$  of (1) on  $I_{-1}$  such that (8) holds.

*Proof.* If  $y(t)$  is a solution of (1) on  $I_{-1}$ , then it is possible to put, e.g.,  $\omega(t) \equiv y(t) - 1$ . Now suppose  $\omega(t)$  is a solution of inequality (7) on  $I_{-1}$ . Let  $\varphi(t)$  be an initial function on  $[t_{-1}, t_0]$  such that  $\varphi(t_0) = \omega(t_0)$  and  $\varphi(t) < \omega(t)$  on  $[t_{-1}, t_0)$ . Prove that  $y^\varphi(t) > \omega(t)$  on  $(t_0, \infty)$ . Suppose the contrary. Then there exists a point  $t^* \in (t_0, \infty)$  such that  $y^\varphi(t) > \omega(t)$  on  $(t_0, t^*)$  and  $y^\varphi(t^*) = \omega(t^*)$ . Note that the case  $y^\varphi(t) \equiv \omega(t)$  on  $[t_0, t^*)$  is impossible because in view of (7) for function  $W(t) \equiv y^\varphi(t) - \omega(t)$  we have

$$\begin{aligned} \left. \frac{dW(t)}{dt} \right|_{t=t_0} &= [\beta(t)[y^\varphi(t) - y^\varphi(t - \tau(t))] - \dot{\omega}(t)]|_{t=t_0} \\ &> \beta(t_0)[\omega(t_0) - \omega(t_0 - \tau(t_0))] - \dot{\omega}(t_0) \geq 0. \end{aligned}$$

Since  $W(t_0) = W(t^*) = 0$ , then (in view of Rolle's theorem), there exists a point  $t^{**} \in (t_0, t^*)$  such that  $W'(t) > 0$ ,  $t \in [t_0, t^*)$  and  $W'(t^{**}) = 0$ . This is impossible. Indeed, let a function  $W_1(t) = W(t) - W(t^{**})$  be involved. Then  $W_1'(t^{**}) = W'(t^{**}) = 0$  and  $y^\varphi(t) < \omega(t) + W(t^{**})$  on  $[t_{-1}, t^{**})$ . However, in view of (7),

$$\begin{aligned} \left. \frac{dW_1(t)}{dt} \right|_{t=t^{**}} &= [\beta(t)[y^\varphi(t) - y^\varphi(t - \tau(t))] - \dot{\omega}(t)]|_{t=t^{**}} \\ &> \beta(t^{**})[\omega(t^{**}) + W(t^{**}) \\ &\quad - \omega(t^{**} - \tau(t^{**})) - W(t^{**})] \\ &\quad - \dot{\omega}(t^{**}) \geq 0. \end{aligned}$$

This contradiction proves the theorem because (8) is valid, e.g., for  $y(t) = y^\varphi(t) + 2 \max_{[t_{-1}, t_0]}(\omega(t) - \varphi(t))$ . From Theorem 1 immediately follows

**THEOREM 2.** Equation (1) has a solution  $y^\varphi(t)$  with property  $y^\varphi(\infty) = \pm \infty$  iff inequality (7) has a solution  $\omega(t)$  with property  $\omega(\infty) = \pm \infty$ .

**Remark 1.** Inequality (7) has a solution  $\omega = \omega(t)$  on  $I_{-1}$  of the form,

(i)

$$\omega(t) = k \int_{t_{-1}}^t \beta(s) ds + p, \quad k \in \mathbb{R}^+, p \in \mathbb{R},$$

if

$$\int_{t-\tau(t)}^t \beta(s) ds \geq 1, \quad t \in I;$$

(ii)

$$\omega(t) = k \exp \left[ \int_{t-1}^t \varepsilon(s) \beta(s) ds \right] + p, \quad k \in \mathbb{R}^+, p \in \mathbb{R},$$

if there is a function  $\varepsilon(t) \in C(I_{-1}, \mathbb{R})$  such that

$$\varepsilon(t) \leq 1 - e^{-\int_{t-1}^t \varepsilon(s) \beta(s) ds}, \quad t \in I;$$

(iii)

$$\omega(t) = k \int_{t-1}^t \left[ \frac{1}{\tau(s)} - \beta(s) \right] ds + p, \quad k \in \mathbb{R}^+, p \in \mathbb{R},$$

if

$$\frac{1}{\tau(t) \beta(t)} - 1 \leq \int_{t-\tau(t)}^t \left[ \frac{1}{\tau(s)} - \beta(s) \right] ds, \quad t \in I.$$

Solution  $\omega(t)$  given by (i) obviously satisfies condition  $\omega(+\infty) = +\infty$ .  
Solution  $\omega(t)$  given by (ii) satisfies condition  $\omega(+\infty) = +\infty$  if, moreover,

$$\int^{+\infty} \varepsilon(s) \beta(s) ds = +\infty,$$

and in the case (iii) for  $\omega(+\infty) = +\infty$  it is sufficient that

$$\int^{+\infty} \left[ \frac{1}{\tau(s)} - \beta(s) \right] ds = +\infty.$$

**THEOREM 3.** For each  $C_1 \in \mathbb{R}$ ,  $C_2 \in \mathbb{R}$ ,  $C_1 < C_2$  there is a nonconstant solution  $y = y(t)$  of (1) on  $I_{-1}$  such that

$$C_1 < y(t) < C_2. \quad (9)$$

*Proof.* Let us show that the set  $\omega = \{(t, y) \in \omega_0: m_1(t, y) < 0, l_1(t, y) < 0\}$  where  $\omega_0 = (t_\varepsilon, \infty) \times \mathbb{R}$ ,  $t_\varepsilon = t_0 - \varepsilon - \tau(t_0 - \varepsilon)$ ,  $\varepsilon \in \mathbb{R}^+$ ,  $t_{-1} + \varepsilon < t_0$ ,  $m_1(t, y) = t_\varepsilon - t$  and  $l_1(t, y) \equiv l_1(y) = (y - C_1)(y - C_2)$  is a regular polyfacial set with respect to (1). The condition  $(\alpha)$  of Definition 1 is

obviously fulfilled. Further  $Dm_1 = -1 < 0$  and

$$Dl_1 = \beta(t)[2\pi(0) - C_1 - C_2][\pi(0) - \pi(-\tau(t))], \quad t > t_0 - \varepsilon.$$

Since  $C_1 < \pi(\theta) < C_2$  where  $\theta \in [-\tau, 0)$ , then

$$Dl_1|_{\pi(0)=C_1} = \beta(t)(C_1 - C_2)(C_1 - \pi(-\tau(t))) > 0, \quad t > t_0 - \varepsilon,$$

and

$$Dl_1|_{\pi(0)=C_2} = \beta(t)(C_2 - C_1)(C_2 - \pi(-\tau(t))) > 0, \quad t > t_0 - \varepsilon.$$

Now let us put  $Z = \{(t, y) \in \bar{\omega}, t = t_0\}$ ,  $\Pi: W \ni (t^*, y^*) \mapsto (t_0, y^*) \in Z \cap W$  and choose a system of initial functions  $p_{Z, \omega}$  which consists of *nonconstant* functions only. (This possibility follows from geometrical considerations.) All assumptions of Lemma 1 hold and then there is a point  $z_0 = (t_0, y_0) \in Z \cap \omega$  such that the graph of the (nonconstant) solution which is defined by function  $p(z_0) \in p_{Z, \omega}$  lies in  $\omega$  for  $t \in D_{t_0, p(z_0)} = [t_0, +\infty)$ , that is its coordinates satisfy the inequalities (9). The theorem is proved.

**THEOREM 4.** *Let  $Y(t)$  be a solution of (1) on  $I_{-1}$  with property  $Y(+\infty) = +\infty$ . Then for each solution  $y(t)$  of (1) on  $I_{-1}$ ,*

$$y(t) = K \cdot Y(t) + \delta(t) \quad (10)$$

*holds, where  $K \in \mathbb{R}$  is a constant, depending on  $y(t)$ , and  $\delta(t)$  is a bounded solution of (1) on  $I_{-1}$  dependent on  $y(t)$ . This representation is unique (with respect to  $K$  and  $\delta(t)$ ).*

*Proof.* Let  $y(t)$  be bounded on  $I_{-1}$ . Then (10) holds for  $K = 0$  and  $\delta(t) \equiv y(t)$  only. Let  $y(t)$  be not bounded. Without loss of generality, we can suppose that both  $Y(t)$  and  $y(t)$  are monotone on  $[t_{-1}, t_0]$  (in opposite case both solutions become monotone in view of Lemma 3 on interval  $[t_k, t_{k+1}]$  where  $t_k = t_{k+1} - \tau(t_{k+1})$ ,  $k \geq 0$  and the reasonings can be made on this interval),  $y(t)$  is monotone decreasing and  $Y(t) > y(t)$  on  $[t_{-1}, t_0]$  (in view of Lemmas 2 and 3). Let  $\varepsilon \in [0, 1]$  be a parameter. Consider the one-parametric family of functions,

$$\Phi(t, \varepsilon) = \varepsilon Y(t) + (1 - \varepsilon)y(t).$$

Obviously  $\Phi(t, 1) = Y(t)$ ,  $\Phi(t, 0) = y(t)$ . In the proof of Theorem 3 we put the system  $\Phi(t, \varepsilon)$ ,  $\varepsilon \in [0, 1]$ ,  $t \in [t_{-1}, t_0]$  as the system of initial functions  $p_{Z, \omega}$ . Put, moreover,  $C_1 = y(t_0)$ ,  $C_2 = Y(t_0)$ ,  $C_1 < C_2$ . By Theorem 3 there is a bounded solution  $\tilde{\delta}(t)$  of (1) such that  $C_1 < \tilde{\delta}(t) < C_2$  on  $I_{-1}$ . This solution corresponds to an initial function for a value of parameter

$\varepsilon = \tilde{\varepsilon} \in (0, 1)$ . Therefore,

$$\tilde{\delta}(t) = \tilde{\varepsilon}Y(t) + (1 - \tilde{\varepsilon})y(t), \quad t \in I_{-1},$$

or

$$y(t) = \frac{\tilde{\varepsilon}Y(t)}{\tilde{\varepsilon} - 1} + \frac{1}{1 - \tilde{\varepsilon}}\tilde{\delta}(t), \quad t \in I_{-1}.$$

Putting  $K = \tilde{\varepsilon}/(\tilde{\varepsilon} - 1)$ ,  $\delta(t) = \tilde{\delta}(t)/(1 - \tilde{\varepsilon})$ , we obtain (10). If this representation is not unique, then for a  $\varepsilon^\diamond \in (0, 1)$ ,  $\varepsilon^\diamond \neq \tilde{\varepsilon}$ ,

$$y(t) = \frac{\varepsilon^\diamond Y(t)}{\varepsilon^\diamond - 1} + \frac{1}{1 - \varepsilon^\diamond}\delta^\diamond(t), \quad t \in I_{-1},$$

where  $\delta^\diamond(t)$  is a solution of (1) bounded on  $I_{-1}$  and

$$\left( \frac{\tilde{\varepsilon}}{\tilde{\varepsilon} - 1} - \frac{\varepsilon^\diamond}{\varepsilon^\diamond - 1} \right) Y(t) = \frac{1}{1 - \varepsilon^\diamond}\delta^\diamond(t) - \frac{1}{1 - \tilde{\varepsilon}}\tilde{\delta}(t).$$

This is impossible because  $\tilde{\varepsilon}(\tilde{\varepsilon} - 1)^{-1} \neq \varepsilon^\diamond(\varepsilon^\diamond - 1)^{-1}$  and  $Y(+\infty) = +\infty$ . The theorem is proved.

*Remark 2.* Obviously, for each  $K \in \mathbb{R}$  and for each bounded solution  $\delta(t)$  of (1) there is a solution  $y(t)$  of (1) given by (10), i.e., the converse affirmation is valid. If Theorem 4 holds, then in the representation (10) the bounded solution  $\delta(t)$  can be written in the form  $\delta(t) = L + \delta_1(t)$  where  $\delta_1(t)$  is a bounded solution of (1),  $L \in \mathbb{R}$  and  $\delta_1(t) = O(L)$ . Then

$$y(t) = K \cdot Y(t) + L + \delta_1(t).$$

This representation is not unique.

## 5. EXPONENTIAL ASYMPTOTICS OF SOLUTIONS

LEMMA 5. If  $y(t)$  is a positive solution of (1) on  $I_{-1}$ , then the expression,

$$V(t) = e^{-\int_{I_{-1}}^t \beta(s) ds} \cdot y(t)$$

is a decreasing function on  $I$ .

*Proof.* For  $V'(t)$  we have

$$\begin{aligned} V'(t) &= e^{-\int_{I_{-1}}^t \beta(s) ds} [y'(t) - \beta(t)y(t)] \\ &= -e^{-\int_{I_{-1}}^t \beta(s) ds} \beta(t)y(t - \tau(t)) < 0, \quad t \in I. \end{aligned}$$

The lemma is proved.



THEOREM 5. *Solution  $y = y(t)$ ,  $t \in I_{-1}$  of (1) is represented in the form,*

$$y(t) = (\xi + \gamma(t))e^{\int_{t-1}^t \beta(s) ds}, \quad t \in I_{-1}, \quad (11)$$

where  $\xi$  is a positive constant and  $\gamma(t)$  is a continuous positive function on  $I$  with property  $\gamma(+\infty) = 0$  iff there is a function  $\alpha(t) \in C(I_{-1}, \mathbb{R}^+)$  such that  $\int^{+\infty} \alpha(s)\beta(s) ds < \infty$  and

$$\alpha(t) \geq e^{-\int_{t-\tau(t)}^t (1-\alpha(s))\beta(s) ds}, \quad t \in I. \quad (12)$$

*Proof.* A. By Theorem 1 and part (ii) of Remark 1, if  $\varepsilon(t) \equiv 1 - \alpha(t)$  is put, there is a solution  $y = y(t)$  of (1) such that

$$y(t) > \omega(t) = ke^{\int_{t-1}^t (1-\alpha(s))\beta(s) ds}, \quad k \in \mathbb{R}^+, t \in I_{-1}. \quad (13)$$

By Lemma 5 there exists  $\lim_{t \rightarrow \infty} V(t) = \xi \geq 0$ , i.e.,

$$y(t) = (\xi + \gamma(t))e^{\int_{t-1}^t \beta(s) ds}, \quad t \in I_{-1}, \quad (14)$$

where  $\gamma(t)$  has properties indicated in formulation of theorem. Comparing (13) and (14) and supposing that  $\xi = 0$ , we have

$$\gamma(t)e^{\int_{t-1}^t \beta(s) ds} > ke^{\int_{t-1}^t (1-\alpha(s))\beta(s) ds},$$

or

$$\gamma(t) > ke^{-\int_{t-1}^t \alpha(s)\beta(s) ds}.$$

The last inequality cannot be valid because the limit of the right-hand side is nonzero for  $t \rightarrow \infty$ . Therefore  $\xi \geq k > 0$ .

B. Let  $y(t)$  be of the form (11). Denote  $\alpha_1(t) \equiv y(t - \tau(t))y^{-1}(t)$ . From (1) we obtain

$$-\frac{\dot{y}(t)}{y(t)} = -\beta(t)(1 - \alpha_1(t)), \quad t \in I,$$

and by integration from  $t - \tau(t)$  to  $t$  we have

$$\alpha_1(t) \equiv e^{\int_{t-\tau(t)}^t (1-\alpha_1(s))\beta(s) ds}, \quad t \in I_1 = [t_1, \infty),$$

that is (12) holds for  $\alpha \equiv \alpha_1(t)$  on  $I_1$ . Moreover, from (1) follows

$$\int_{t_0}^t \left[ \frac{\dot{y}(s)}{y(s)} - \beta(s) \right] ds = -\int_{t_0}^t \alpha_1(s)\beta(s) ds,$$

and, using (11), we have

$$\frac{\xi + \gamma(t)}{\xi + \gamma(t_0)} = e^{-\int_{t_0}^t \alpha_1(s) \beta(s) ds}.$$

From this follows the convergence of integral  $\int^{+\infty} \alpha_1(s) \beta(s) ds$ . Defining  $\alpha(t - \tau(t)) \equiv \alpha_1(t)$  in (12) means the end of the proof.

*Remark 3.* The function  $\alpha(t)$  often can be taken in the form  $\alpha(t) = [\beta^{p+1}(t)]^{-1}$ ,  $p \in \mathbb{R}^+$  if  $\int^\infty \beta^{-p}(s) ds < \infty$ . E.g., if  $\beta(t) = t$  and  $\tau(t) \in [\tau_1, \tilde{\tau}]$ ,  $0 < \tau_1 = \text{const}$ , then it is possible to put (if  $t_{-1}$  is sufficiently large)  $\alpha(t) = t^{-3}$ .

**THEOREM 6.** *Let there be a solution  $y(t)$  of (1) in the form (11) on  $I_{-1}$  and*

$$\int^\infty \beta(s) e^{-\int_{s-\tau(s)}^s \beta(q) dq} ds < \infty.$$

*Then*

$$\gamma(t) > \left[ \exp \left( \int_t^\infty \beta(s) e^{-\int_{s-\tau(s)}^s \beta(q) dq} ds \right) - 1 \right] \cdot \xi, \quad t \in I_1. \quad (15)$$

*Proof.* From Lemma 5 we have  $V(t) < V(t - \tau(t))$ ,  $t \in I_1$  and

$$V'(t) = -\beta(t) e^{-\int_{t-1}^t \beta(s) ds} y(t - \tau(t)) < -\beta(t) e^{-\int_{t-\tau(t)}^t \beta(s) ds} V(t). \quad (16)$$

Since  $V(\infty) = \xi > 0$  (the constant  $\xi$  is the same as in Theorem 5) then, by integration (16) from  $t$  to  $\infty$  we have

$$V(t) = \xi + \gamma(t) > \xi \exp \left( \int_t^\infty \beta(s) e^{-\int_{s-\tau(s)}^s \beta(q) dq} ds \right), \quad t \in I_1.$$

From this follows (15). The theorem is proved.

**COROLLARY 1.** *Let Theorem 5 hold, let a solution  $y = y_1(t)$  of (1) with representation (11) be fixed and  $\int^\infty \beta(s) ds = \infty$ . Then (as it follows from Theorem 4) for each solution  $y(t)$  of (1) on  $I_{-1}$  there is a constant  $K \in \mathbb{R}$  and a bounded solution  $\delta(t)$  of (1) such that*

$$y(t) = Ky_1(t) + \delta(t) = K(\xi + \gamma(t)) e^{\int_{t_0}^t \beta(s) ds} + \delta(t), \quad t \in I_{-1}. \quad (17)$$

*If, moreover, Theorem 6 holds, then (15) is valid.*

## 6. OTHER ASYMPTOTICS

**THEOREM 7.** *The existence of functions  $\varphi_i \in C([t_\varepsilon, \infty), \mathbb{R}) \cap C^1([t_0 - \varepsilon), \mathbb{R})$ ,  $i = 1, 2$ , where  $\varepsilon \in \mathbb{R}^+$ ,  $\varepsilon = \text{const}$ ,  $t_{-1} + \varepsilon < t_0$ ,  $t_\varepsilon = t_0 - \varepsilon - \tau(t_0 - \varepsilon)$ ,  $\varphi_2(t) \leq \varphi_1(t)$ ,  $t \in I_{-1}$  and*

$$\begin{aligned}\varphi_1'(t) &\leq \beta(t)[\varphi_1(t) - \varphi_1(t - \tau(t))], & t > t_0 - \varepsilon, \\ \varphi_2'(t) &\geq \beta(t)[\varphi_2(t) - \varphi_2(t - \tau(t))], & t > t_0 - \varepsilon\end{aligned}$$

*is sufficient and necessary for existence of solution  $y = y(t)$  of (1) such that*

$$\varphi_2(t) \leq y(t) \leq \varphi_1(t), \quad t \in I_{-1}. \quad (18)$$

*Proof.* Necessity is obvious, because  $\varphi_1(t) \equiv \varphi_2(t) \equiv y(t)$  can be taken.

*Sufficiency.* Let us choose an  $\delta \geq 0$  such that  $\varphi_2(t) < \tilde{\varphi}_1(t) \equiv \varphi_1(t) + \delta$  on  $I_{-1}$ . Using topological principle (as in proof of Theorem 3) putting  $\omega = \{(t, y) \in \omega_0: m_1(t, y) < 0, l_1(t, y) < 0\}$  where  $\omega_0 = (t_\varepsilon, \infty) \times \mathbb{R}$ ,  $m_1(t, y) = t_\varepsilon - t$  and  $l_1(t, y) \equiv l_1(y) = (y - \tilde{\varphi}_1(t))(y - \varphi_2(t))$ , we can prove that there exists a solution  $y = y_\delta(t)$  on  $I_{-1}$  (generated, e.g., by an initial function  $\Phi_\delta(t, \varepsilon) = \varepsilon(\varphi_1(t) + \delta) + (1 - \varepsilon)\varphi_2(t)$ ,  $t \in [t_{-1}, t_0]$  where  $\varepsilon \in (0, 1)$  is a fixed number not depending on  $\delta$  in further reasonings) which satisfies inequalities,

$$\varphi_2(t) < y_\delta(t) < \tilde{\varphi}_1(t), \quad t \in I_{-1}.$$

Taking a sequence  $\{\delta_n\} \rightarrow 0^+$ , we conclude that the limit solution  $y_0(t) = \lim_{n \rightarrow \infty} y_{\delta_n}(t)$ , generated by initial function  $\Phi_0(t, \varepsilon)$ , exists and satisfies (18). If the left inequality of (18) does not hold, then there is  $t^* \in I$  such that  $y_0(t^*) < \varphi_2(t^*)$ . However, this is impossible in view of continuous dependence of solutions on initial data. The case  $y_0(t^{**}) > \varphi_1(t^{**})$  for a  $t^{**} \in I$  is impossible in view of construction. The theorem is proved. From Theorems 4 and 7 follows

**THEOREM 8.** *Let there be functions  $\varphi_i(t)$ ,  $i = 1, 2$ , satisfying all conditions described in Theorem 7 and, moreover,  $\varphi_1(\infty) = +\infty$ ,  $\lim_{t \rightarrow \infty} \varphi_1(t)(\varphi_2(t))^{-1} = 1$ . Then each solution  $y = y(t)$  of (1) can be expressed in the form,*

$$y(t) = K\varphi_i(t)[1 + (-1)^i \tilde{\gamma}_i(t)] + \tilde{\delta}_i(t), \quad t \in I_{-1}, i = 1, 2, \quad (19)$$

*where  $K \in \mathbb{R}$  is a constant dependent on  $y(t)$ ;  $\tilde{\gamma}_i(t) \geq 0$  and  $\tilde{\delta}_i(t)$  are continuous bounded functions dependent on  $y(t)$  and  $\tilde{\gamma}_i(\infty) = 0$ .*

EXAMPLE 1. Consider the equation,

$$\dot{y}(t) = \left( \frac{1}{\tau} + \frac{p-1}{2t} \right) [y(t) - y(t-\tau)], \quad (20)$$

where  $p > 1$ ,  $\tau > 0$ ,  $\tau = \text{const}$ . By Theorem 8 (suppose that  $t_0$  is sufficiently large) each solution of (20) is representable in the form (19) where  $\varphi_1(t) = (t+C)^p$ ,  $C = \text{const}$ ,  $C > \tau(p+1)/6$ ,  $\varphi_2(t) = t^p$ . E.g., for  $i = 2$  we have

$$y(t) = Kt^p[1 + \tilde{\gamma}_2(t)] + \tilde{\delta}_2(t), \quad t \in I_{-1},$$

if  $t_0$  is sufficiently large.

From Theorems 7 and 8 it follows

THEOREM 9. Let  $\tau(t) \equiv \tau \in \mathbb{R}^+$ ,  $\beta \in C([t_{-1} - \tau + b, \infty), \mathbb{R}^+)$ ,  $a + \tau > b \geq 0$ ,  $a, b \in \mathbb{R}$  and

$$\beta(t-b) \leq \frac{\beta(t)}{\int_{t-\tau}^t \beta(s) ds} \leq \beta(t-a-\tau), \quad t > t_{-1} + b.$$

If, moreover,  $\lim_{t \rightarrow \infty} \beta(t-b)/\beta(t-a-\tau) = 1$  and  $\int^\infty \beta(s) ds = \infty$ , then each solution  $y = y(t)$  of (1) can be expressed in the form,

$$y(t) = K\varphi_i(t)[1 + (-1)^i \tilde{\gamma}_i(t)] + \tilde{\delta}_i(t), \quad t \in I_{-1}, i = 1, 2,$$

where  $\varphi_2(t) = \int_{t_0-\tau}^{t+b} \beta(s) ds$ ,  $\varphi_1(t) = \int_{t_0-\tau}^{t+a+\tau} \beta(s) ds$ , and  $K \in \mathbb{R}$  is a constant dependent on  $y(t)$ ;  $\tilde{\gamma}_i(t) \geq 0$  and  $\tilde{\delta}_i(t)$  are continuous bounded functions dependent on  $y(t)$  and  $\tilde{\gamma}_i(\infty) = 0$ .

EXAMPLE 2. Conditions of Theorem 9 are satisfied, e.g., if  $\beta(t) = \tau^{-1}(1 + \varepsilon \sin(2\pi/\tau)t)$  where  $\varepsilon = \text{const}$ ,  $|\varepsilon| < 1$ , and  $a = b = 0$ . In this case we conclude that each solution  $y = y(t)$  of (1) can be expressed in the form,

$$y(t) = Kt[1 + (-1)^i \tilde{\gamma}_i(t)] + \tilde{\delta}_i(t), \quad t \in I_{-1}, i = 1, 2,$$

where  $K \in \mathbb{R}$  is a constant dependent on  $y(t)$ ,  $\tilde{\gamma}_i(t) \geq 0$ , and  $\tilde{\delta}_i(t)$  are continuous functions dependent on  $y(t)$  and  $\tilde{\gamma}_i(\infty) = 0$ .

## 7. APPLICATIONS AND COMPARISONS

(A) In the paper S. N. Zhang [18] there is proved, e.g., the result about asymptotic structure of solutions of (1) under main assumptions  $\beta(t) > 0$ ,

$t > 0$ ,  $\tau(t) \equiv 1$  and

$$\int_t^{t+1} \beta(s) ds \geq 1, \quad \int_t^{t+1} \beta(s) ds \neq 1, \quad (21)$$

for  $t \geq T_0 > 0$ . These conditions are a partial case of condition (i) in Remark 1. Consider, moreover, the equation,

$$\dot{y}(t) = \left( \frac{1}{\tau} - \frac{a}{t} \right) [y(t) - y(t - \tau)], \quad (22)$$

where  $\beta(t) = (\tau)^{-1} - at^{-1}$ ,  $\tau(t) \equiv \tau = \text{const}$ ,  $a = \text{const}$ , and  $a \in (0, \frac{1}{2}]$ . Then neither conditions (21) for  $\tau = 1$  nor condition (i) in Remark 1 hold. In this case the condition (iii) in Remark 1 holds (if  $t_{-1}$  is sufficiently large),  $\omega(+\infty) = +\infty$  and therefore there exists a nonbounded positive solution  $y(t)$  of (1) which tends to infinity and satisfies inequality  $y(t) > a \ln t / t_{-1}$ . Clearly, Theorem 4 holds. This examples shows simultaneously that the value  $a = \frac{1}{2}$  is a bifurcation value. Indeed, in the paper of F. V. Atkinson and J. R. Haddock [2] it is shown that in the case  $a > \frac{1}{2}$  each solution of (22) tends to a constant.

(B) The equation of the type (2) where  $c \in C(I_{-1}, \mathbb{R}^+)$  can be obtained from equation (1) by transformation  $y(t) = x(t) \exp(\int_{t_0}^t \beta(s) ds)$ . Conversely, if there is a positive solution of (2)  $x(t) = \zeta(t)$  on  $I_{-1}$ , then the transformation  $x(t) = \zeta(t)y(t)$  gives

$$\dot{y}(t) = - \frac{\zeta'(t)}{\zeta(t)} [y(t) - y(t - \tau(t))], \quad t \in I_{-1}, \quad (23)$$

that is (23) has the form of (1) with  $\beta(t) \equiv -\zeta'(t)/\zeta(t)$ . Note that sufficient and necessary conditions for existence of a positive solution of (2) are given by Detang Zhou [19] (see L. H. Erbe, Qingkai Kong, and B. G. Zhang [6] too).

**THEOREM 10.** Suppose  $c \in C([t_\varepsilon, \infty), \mathbb{R}^+)$ , where  $\varepsilon \in \mathbb{R}^+$ ,  $\varepsilon = \text{const}$ ,  $t_{-1} + \varepsilon < t_0$ ,  $t_\varepsilon = t_0 - \varepsilon - \tau(t_0 - \varepsilon)$ ,  $c(t) < 1$  on  $I_{-1}$ . Let, moreover, there be a positive constant  $L$  such that

$$c(t) \leq L \left( \ln \frac{1}{c(t)} \right) \exp \left( L \int_{t-\tau(t)}^t \ln c(s) ds \right), \quad t > t_0 - \varepsilon.$$

Then there is a positive solution of (2)  $x = \zeta(t)$  such that

$$0 < \zeta(t) < \exp \left( L \int_{t_0}^t \ln c(s) ds \right), \quad t \in I_{-1}. \quad (24)$$

*Proof.* The existence of a positive solution of (2)  $x = \zeta(t)$  with property (24) can be proved by means of topological principle analogously as in the proof of Theorem 3. For this it is necessary to choose  $\omega = \{(t, y) \in \omega_0: m_1(t, y) < 0, l_1(t, y) < 0\}$  where  $\omega_0 = (t_\varepsilon, \infty) \times \mathbb{R}$ ,  $m_1(t, y) = t_\varepsilon - t$ , and  $l_1(t, x) \equiv l_1(x) = x \cdot (x - \exp(L \int_{t_0}^t \ln c(s) ds))$ . The details are omitted.

**THEOREM 11.** *Let Theorem 10 hold. Suppose, moreover, that there is a constant  $p \in (0, 1]$  such that  $\int^\infty c^p(s) ds < \infty$ ,  $\int^\infty \ln c(s) ds = -\infty$  and*

$$c^{p-1}(t) \geq \exp\left(\int_{t-\tau(t)}^t c^p(s) ds\right), \quad t \in I_{-1}. \quad (25)$$

*Then for each solution of (2)  $x = x(t)$  there is a constant  $K \in \mathbb{R}$  and a bounded solution of (23)  $\delta(t)$  such that*

$$x(t) = K\zeta(t_0)(\xi + \gamma(t)) + \zeta(t)\delta(t), \quad t \in I_{-1}, \quad (26)$$

*where  $\zeta(t)$  is the positive solution of (2) given by Theorem 10,  $\xi > 0$  is a fixed constant and  $\gamma(t) \geq 0$  is a fixed continuous function with property  $\gamma(\infty) = 0$ .*

$$\int_{t_0}^\infty \beta(s) ds = - \int_{t_0}^\infty \frac{\zeta'(s)}{\zeta(s)} ds = -\ln \frac{\zeta(+\infty)}{\zeta(t_0)} = +\infty.$$

Put  $\alpha(t) = -c^p(t)\zeta(t)/\zeta'(t)$ . Without loss of generality we suppose that  $\zeta \in C^1(I_{-1}, \mathbb{R}^+)$ . We see that  $\int^\infty \alpha(s)\beta(s) ds = \int^\infty c^p(s) ds < \infty$  and (12) holds in view of (25). Therefore, by Corollary 1 for each solution of (23), there is a constant  $\xi \in \mathbb{R}$  and a bounded function  $\delta(t)$  continuous on  $I_{-1}$  such that representation (17) holds and, consequently, for each solution of (2) there is a constant  $\xi \in \mathbb{R}$  and a bounded function  $\delta(t)$  continuous on  $I_{-1}$  such that

$$x(t) = K\zeta(t_0)(\xi + \gamma(t)) + \zeta(t)\delta(t).$$

The theorem is proved.

The proof of the following theorem is similar to that of Theorem 11 and hence is omitted.

**THEOREM 12.** *Let there be a positive solution of (2)  $x(t) = \zeta(t)$  on  $I_{-1}$  such that  $\zeta(\infty) = 0$  and  $c \in C([t_\varepsilon, \infty), \mathbb{R}^+)$ . Assume, moreover, that there is a constant  $p \in (0, 1]$  such that  $\int^\infty c^p(s) ds < \infty$  and that (25) holds. Then for each solution of (2)  $x = x(t)$  there is a constant  $K \in \mathbb{R}$  and a bounded solution of (23)  $\delta(t)$  such that the representation (26) holds where  $\xi > 0$  is a fixed constant and  $\gamma(t) \geq 0$  is a fixed continuous function with property  $\gamma(\infty) = 0$ .*

*Remark 4.* The existence of a positive solution (without guarantee that its asymptotic behavior is described by inequalities (24)) follows from (25) because previously mentioned sufficient and necessary conditions are valid (see [19, 6]). Note, moreover, that, by means of appropriate transformation of variables, it is possible, under our assumptions, to obtain equations with *constant* delay that are equivalent to (1) and (2). This follows from results given in [13, 14]. If Theorems 11 or 12 hold, then (as it follows from formula (26)) the trivial solution of (2) is *nonasymptotically* stable on  $I_{-1}$  because by Remark 2 there is a solution of (2) for each  $K \neq 0$  and therefore for  $K \neq 0 \lim_{t \rightarrow \infty} x(t) = K\zeta(t_0)\xi \neq 0$ .

EXAMPLE 3. Consider the equation (see, e.g., J. Hale [10]),

$$\dot{x}(t) = -2te^{1-2t}x(t-1).$$

Because it follows from Theorems 11 and 12, each solution of this equation is representable in the form (26). Because there is a positive solution  $\zeta(t) = \exp(-t^2)$ , this representation can be written in the form,

$$x(t) = Ke^{-t_0^2}(\xi + \gamma(t)) + e^{-t^2}\delta(t), \quad t \in I_{-1}.$$

Moreover, by Remark 4, the trivial solution of this equation is nonasymptotically stable.

## ACKNOWLEDGMENT

This work was supported by the grant 201/96/0410 of Czech Grant Agency (Prague).

## REFERENCES

1. O. Arino, I. Györi, and M. Pituk, Asymptotically diagonal delay differential systems, *J. Math. Anal. Appl.* **204** (1996), 701–728.
2. F. V. Atkinson and J. R. Haddock, Criteria for asymptotic constancy of solutions of functional differential equations, *J. Math. Anal. Appl.* **91** (1983), 410–423.
3. R. Bellman and K. L. Cooke, Differential-difference equations, in “Mathematics in Science and Engineering, A Series of Monographs and Textbooks,” Academic Press, New York, London, 1963.
4. J. Diblík, Bounded solutions with positive coordinates of functional-differential equations of the retarded type, in “Proceedings of the Conference on Ordinary Differential Equations,” (P. Marušák, Ed.), Poprad, (Slovak Republic), pp. 13–22, 1994.
5. J. Diblík, Asymptotic behaviour of solutions of linear differential equations with delay, *Ann. Polon. Math.* **LVIII.2** (1993), 131–137.
6. L. H. Erbe, Qingkai Kong, and B. G. Zhang, “Oscillation Theory for Functional Differential Equations,” Marcel Dekker, New York, Basel, Hong Kong, 1995.

7. I. Györi and M. Pituk,  $L^2$ -perturbation of a linear delay differential equation, *J. Math. Anal. Appl.* **195** (1995), 415–427.
8. I. Györi and M. Pituk, Comparison theorems and asymptotic equilibrium for delay differential and difference equations, *Dynamic Systems and Applications* **5** (1996), 277–302.
9. K. Gopalsamy, “Stability and Oscillations in Delay Differential Equations of Population Dynamics,” Kluwer Academic, Dordrecht, Boston, London, 1992.
10. J. Hale, “Theory of Functional Differential Equations,” Springer-Verlag, New York, Heidelberg, Berlin, 1977.
11. E. Kozakiewicz, On the asymptotic behaviour of positive solutions of two different inequalities with retarded argument, *Colloquia Mathematica Societatis János Bolyai*, **15**, *Differential Equations*, Keszthely (Hungary), 1975, 309–319.
12. T. Krisztin, Asymptotic estimation for functional differential equations via Lyapunov functions, Qualitative theory of differential equations, Szeged (Hungary) *Colloq. Math. Soc. János Bolyai*, **53** (1988), 365–376.
13. F. Neuman, On equivalence of linear functional-differential equations, *Results in Mathematics*, **26** (1994), 354–359.
14. F. Neuman, On transformations of differential equations and systems with deviating argument, *Czechoslovak Math. J.* **31** (1981), 87–90.
15. B. S. Razumikhin, “Stability of hereditary systems,” Nauka, Moscow, 1988. [In Russian]
16. K. P. Rybakowski, Ważewski’s principle for retarded functional differential equations, *J. Differential Equations*, **36** (1980), 117–138.
17. T. Ważewski, Sur un principe topologique de l’examen de l’allure asymptotique des intégrales des équations différentielles ordinaires, *Ann. Soc. Polon. Math.* **20** (1947), 279–313.
18. S. N. Zhang, Asymptotic behaviour and structure of solutions for equation  $\dot{x}(t) = p(t)[x(t) - x(t - 1)]$ , *J. Anhui University (Natural Science Edition)* **2** (1981), 11–21. [In Chinese]
19. Detang Zhou, Negative answer to a problem of Györi, *J. Shandong University* **24** (1989), 117–121.